

B.A./B.Sc 4th Semester (Honours) Examination, 2019 (CBCS)

Subject : Mathematics

Paper : BMH4 CC08

Time: 3 Hours

Full Marks: 60

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.*

[Notations and Symbols have their usual meaning.]

Group-A

Marks : 20

1. Answer any ten questions:

2×10=20

(a) Evaluate : $\int_0^3 [x] dx$, where $[x]$ denotes the largest integer not larger than real number x .(b) Let $f(x) = x$ if $x \neq \frac{1}{n}$

$$f\left(\frac{1}{n}\right) = 1 \quad \text{for } n = 1, 2, 3, \dots$$

Is $f(x)$ Riemann integrable in $[0, 1]$? Support your answer.(c) Give examples of two functions f and g which are not Riemann integrable in $[0, 1]$ but $\phi(x) = \text{Max}\{f(x), g(x)\}$ is Riemann integrable in $[0, 1]$. Answer with justification.(d) Find $\lim_{x \rightarrow 0} \frac{1}{x} \int_0^x \cos \frac{1}{t^2} dt$, if it exists.(e) Test the convergence of $\int_0^\infty e^{-x^2} dx$.(f) Let f and g be two continuous functions defined over $[a, b]$, $a < b$, such that

$$\int_a^b f(x) dx = \int_a^b g(x) dx. \text{ Show that there exists } c \in [a, b] \text{ such that } f(c) = g(c).$$

(g) Test for convergence of the integral $\int_0^1 \frac{\sqrt{x}}{\sin x} dx$.(h) With proper justification give an example of a sequence of functions $\{f_n\}$ which converges pointwise to a function f such that each f_n is Riemann integrable but f is not Riemann integrable.(i) Test for uniform convergence of the sequence $\{(\sin x)^n\}$ in the interval $[1, 2]$.(j) Let $f_n(x) = 1$ if $\frac{1}{n} \leq x \leq 1$ and
 $= nx$ if $0 \leq x < \frac{1}{n}$.Find $\lim_{n \rightarrow \infty} f_n(x)$ in $[0, 1]$, if it exists.

- (k) Test the series $\sum_{n=1}^{\infty} x^n (1-x^n)$ for pointwise convergence and uniform convergence in $[0, 1]$.
- (l) Give an example of a power series which converges nowhere except $x = 0$. Justify your answer.
- (m) Find the region of convergence of the power series $\sum_{n=1}^{\infty} \frac{n!(x+2)^n}{n^n}$.
- (n) Examine whether $\sum_{n=1}^{\infty} (\sin nx + \cos nx)$ is a Fourier series for some bounded integrable function over $[-\pi, \pi]$.
- (o) Show that the Fourier series of a bounded integrable odd function over $[-\pi, \pi]$ consists of sine terms only.

Group-B

Marks : 20

2. Answer any four questions from the following:

5×4=20

(a) State and prove Fundamental Theorem of Integral Calculus.

2+3=5

(b) If $\{f_n(x)\}$ is a sequence of continuous functions on $[a, b]$ and if $f_n(x) \rightarrow f(x)$ uniformly in $[a, b]$ as $n \rightarrow \infty$, then prove that $f(x)$ is continuous in $[a, b]$.(c) (i) Prove that every monotone function over $[a, b]$ is Riemann integrable therein.(ii) Give an example with proper justification of a function f on $[0, 1]$ which is not Riemann integrable on $[0, 1]$ but $|f|$ is Riemann integrable on $[0, 1]$.

3+2=5

(d) Suppose $f(x) = x^2$ in $0 \leq x \leq 2\pi$ and $f(x)$ is a periodic function of period $= 2\pi$. Obtain the Fourier series of f and examine the convergence of the series in $[0, 2\pi]$.

3+2=5

(e) State and prove Weierstrass M -test for uniform convergence of a series of functions.

2+3=5

(f) State Dirichlet's condition concerning convergence of Fourier series of a function. If f is bounded and integrable in $[-\pi, \pi]$ and if a_n, b_n are its Fourier co-efficients, then prove that

$$\sum_{n=1}^{\infty} (a_n^2 + b_n^2) \text{ is convergent.}$$

2+3=5

Group-C

Marks : 20

3. Answer any two questions from the following:

10×2=20

(a) (i) Let f be a bounded function defined on the closed interval $[a, b]$. Prove that a necessary and sufficient condition that f be Riemann integrable over $[a, b]$ is that to every $\epsilon > 0$ there corresponds a $\delta > 0$ such that for every partition P of $[a, b]$ with $\|P\| \leq \delta$, the oscillatory sum $W(P, f) < \epsilon$.

(ii) If a bounded function f is Riemann integrable in $[a, b]$, show that $|f|$ is also Riemann integrable in $[a, b]$ and $\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$.

(iii) Show that $\left| \int_a^b \frac{\sin x}{x} dx \right| \leq \frac{4}{a}$ for $0 < a < b < \infty$. 4+4+2=10

(b) (i) Prove the relation $\beta(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$, $m > 0$, $n > 0$.

(ii) Let $f(x) = |x|$ for $-\pi < x \leq \pi$ and $f(x + 2\pi) = f(x)$ for all x . Find the Fourier series of f . Hence deduce that $\frac{\pi^2}{8} = \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2}$. 5+(4+1)=5

(c) (i) Test the convergence of the integral $\int_0^{\infty} e^{-x} x^{n-1} dx$, for different values of n .

(ii) Show that $\frac{1}{2} \{\log(1+x)\}^2 = \frac{1}{2} x^2 - \frac{1}{3} \left(1 + \frac{1}{2}\right) x^3 + \frac{1}{4} \left(1 + \frac{1}{2} + \frac{1}{3}\right) x^4 - \dots \forall x \in (-1, 1)$.

5+5=10

(d) (i) Suppose that a power series $f(x) = \sum_{n=0}^{\infty} a_n x^n$ has a finite non-zero radius of convergence ρ . Prove that the series converges uniformly in each compact subset of $(-\rho, \rho)$.

(ii) If two power series $\sum_{n=0}^{\infty} a_n x^n$ and $\sum_{n=0}^{\infty} b_n x^n$ converge to the same sum function in an interval $(-r, r)$, $r > 0$, then show that $a_n = b_n$ for all n .

(iii) Find the radius of convergence of the power series $\sum_{n=1}^{\infty} a_n x^n$, where $a_{2n} = \frac{1}{3^n}$, $a_{2n-1} = \frac{1}{3^{n+1}}$.

4+3+3=10