

**B.Sc. 5th Semester (Honours) Examination, 2022 (CBCS)****Subject : Physics****Course : CC-XI****Quantum Mechanics & Applications****Time: 2 Hours****Full Marks: 40***The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.**Symbols and abbreviations have their usual meanings.***1. Answer any five of the following questions:****2×5=10**

(a) The trial wavefunction of the one-dimensional infinite square well is given by

$$\Psi(x) = \begin{cases} Ax & \text{if } 0 \leq x \leq \frac{a}{2} \\ A(a-x) & \text{if } \frac{a}{2} \leq x \leq a \\ 0 & \text{otherwise} \end{cases}$$

sketch  $\Psi(x)$  and determine the normalization constant  $A$ .

- (b) Consider a system described by a state, which is a superposition of two orthonormal states  $\Phi_1$  and  $\Phi_2$  as  $\Psi = \frac{\sqrt{3}}{2}\Phi_1 + \frac{1}{2}\Phi_2$ . Also consider an ensemble of 50 identical systems each one of them in the state of  $\Psi$ . If measurements are done on all of them, how many systems will be found in each of the states  $\Phi_1$  and  $\Phi_2$ ?
- (c) Find the spectroscopic notation for the ground state configuration of Aluminum  $Al(Z = 13)$  and Scandium ( $Z = 21$ ).
- (d) Differentiate between a weak-field Zeeman effect and the Paschen-Back effect in terms of the relative energy level spacings due to the spin-orbit effect and Zeeman effect.
- (e) Which among the following wave functions represents physically acceptable functions:  
 $f(x) = 3\sin\pi x$ ,  $g(x) = 4 - |x|$ ,  $h^2(x) = 5x$ .
- (f) What is Bohr magneton? Calculate its value.
- (g) Why do all alkali atoms have qualitatively similar spectra?

- (h) Consider a system of three non-interacting identical spin  $\frac{1}{2}$  particle (each of mass  $m$ ) that are in the same spin state  $\left|\frac{1}{2}, \frac{1}{2}\right\rangle$  and confined in one-dimensional infinite well of length  $a$ ;  $V(x) = 0$  for  $0 < x < a$  &  $V(x) = \infty$  for other values of  $x$ . (Assume  $E_n = \frac{n^2\pi^2\hbar^2}{2ma^2}$ ). Determine energy of the ground state and the first excited state of the system.

2. Answer any two of the following questions:

5×2=10

- (a) Given that  $\widehat{p}_r = -i\hbar\left(\frac{\partial}{\partial r} + \frac{1}{r}\right)$ . Find the uncertainty  $\Delta p_r$  in the ground state,

$$\Psi_0(r) = \frac{1}{\sqrt{\pi a_0^3}} e^{-\frac{r}{a_0}}.$$

- (b) (i) Show that the momentum operator  $\widehat{p}_x$  is hermitian.  
 (ii) Show that the commutator  $[\widehat{L}_z, \widehat{L}_-]$  is equal to  $-\hbar\widehat{L}_-$ , where all the symbols have their usual meaning.

2+3=5

- (c) (i) Write a qualitative explanation of the fine-structure splitting of spectral lines in terms of the interaction of spin with magnetic moment.

- (ii) The yellow line of sodium atom is split into two lines having wavelengths 589.0 nm and 589.6 nm because of the transitions  $3P_{\frac{3}{2}} \rightarrow 3S_{\frac{1}{2}}$  and  $3P_{\frac{1}{2}} \rightarrow 3S_{\frac{1}{2}}$  respectively. Find the value of the effective magnetic field on the outer electron, which causes this splitting. [Express in terms of Bohr Magnetron.]

2+3=5

- (d) (i) Write the statistical interpretation of the wave function  $\Psi(x, t)$ .

- (ii) Show with the help of the Schrödinger equation that the space-integrated probability is independent of time. What is the significance of this result so far as the normalization of the wave function is concerned.

1+4=5

3. Answer any two of the following questions:

10×2=20

- (a) (i) Write down Schrödinger equation for the electron of hydrogen atom assuming the nucleus to be stationary. By separation of the variables, obtain the radial equation.

- (ii) The normalized wave function of the ground state of the hydrogen atom is given by

$$\Psi(r) = \frac{1}{\sqrt{\pi a_0^3}} e^{-\frac{r}{a_0}}$$

Find the distance from the nucleus at which the electron is most likely to be found.

- (iii) Argue from uncertainty principle that only one component of the angular momentum of the electron should be quantized.

5+3+2=10

- (b) (i) Write down the Schrödinger equation for a linear harmonic oscillator having mass  $m$ , force constant  $k$  and frequency  $\nu$ .

- (ii) Obtain the energy eigenvalues and eigenfunctions of the oscillator using the Frobenius method. 1+9=10
- (c) (i) Using vector model determine the possible terms corresponding to the principal quantum numbers  $n = 3$  and compute the angle between  $\vec{l}$  and  $\vec{s}$  vectors for the term  $2D_{\frac{5}{2}}$ .

- (ii) Given that the energy of a hydrogen like atom taking into account fine structure corrections is of the form

$$E_{nl} = E_n \left[ 1 + \frac{Z^2 \alpha^2}{n^2} \left( \frac{n}{l + \frac{1}{2}} - \frac{3}{4} \right) \right]$$

where  $n$  and  $l$  represent the principal and orbital angular momentum quantum number respectively and  $\alpha = \frac{1}{137}$ . Show that for fixed  $n$  the difference between the maximum and minimum energy as  $l$  varies from 0 to  $n - 1$  is equal to  $E_n \cdot \frac{Z^2 \alpha^2 4(n-1)}{n(2n-1)}$ . 6+4=10

- (d) (i) What was the purpose of the Stern-Gerlach experiment? Why is a non-uniform magnetic field used in this experiment? In a Stern-Gerlach experiment, always a beam of neutral atoms is used and not ions. — Explain why.
- (ii) Elucidate the space quantization of electron spin and describe its demonstration by the Stern-Gerlach experiment. (1+1+1)+(2+5)=10