

**B.Sc. 5th Semester (Honours) Examination, 2023 (CBCS)****Subject : Physics****Course : CC-XI****(Quantum Mechanics & Applications)****Time: 2 Hours****Full Marks: 40***The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.***1. Answer any five of the following questions:**

2×5=10

- (a) The wavefunction of a particle in a stationary state with an energy  $E_0$  at the time  $t = 0$  is  $\psi(x)$ . After how much time will the wavefunction again be  $\psi(x)$ ?
- (b) Find the value of the constant  $A$  that makes  $e^{-ax^2}$  an eigenfunction of the operator  $\left(\frac{d^2}{dx^2} - Ax^2\right)$ . What is the corresponding eigenvalue? 1+1
- (c) The  $2p$  state for the hydrogen atom is known to be  $re^{-r/2a_0}(\cos\theta)$ . Find out the expectation value of  $r$  in this state.
- (d) Prove that the energy eigenfunction of a free particle is doubly degenerate.
- (e) Explain why normal Zeeman effect occurs only in atoms with even number of electrons.
- (f) Find the value of the Lande g-factor for energy level  $^3P_1$ .
- (g) Find the eigenfunctions of the angular momentum operator  $L_z = i\hbar \frac{\partial}{\partial \phi}$ .
- (h) Evaluate the following commutator  $[\vec{L}, \vec{S}, \vec{J}^2]$ .

**2. Answer any two of the following questions:**

5×2=10

- (a) Using Heisenberg's uncertainty principle, find the first Bohr radius of Hydrogen atom assuming proton to be at rest.
- (b) Consider an electron impinging on a rectangular potential barrier of height  $V_0 = 5\text{eV}$  and thickness  $a = 10^{-10}\text{m}$ . If the kinetic energy of the electron is  $2.5\text{eV}$ , calculate the transmission coefficient.
- (c) What is Zeeman effect? Describe the experimental arrangement for studying the Zeeman effect. 1+4
- (d) Write down the Schrödinger equation for the electron of tritium ( $\text{H}_3$ ) atom, assuming the nucleus to be stationary. Obtain the radial equation by separation of variables with special emphasis on effective potentials.

3. Answer any two of the following questions:

10×2=20

- (a) (i) The wavefunction corresponding to the first excited state of a harmonic oscillator of frequency  $\omega_0$  is given by  $\psi(x) = Ax e^{-\alpha x^2/2}$ ;  $\alpha = \frac{m\omega_0}{\hbar}$ . Sketch  $\psi(x)$  and determine A.
- (ii) Find the expectation value of the operator  $\hat{x}\hat{p}_x$  in this state.
- (iii) An electron of energy 342eV is confined in a one-dimensional box of length 1Å. Find out the quantum number of the electron and energy needed to excite it in the next higher level. (1+2)+4+3

- (b) Write down the time-independent Schrödinger equation for the motion of the electron in a hydrogen atom, assuming that the proton is at rest.

$$\text{Given: } \nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2}$$

Separate the Schrödinger equation into one radial and two angular parts.

1+3+3+3

- (c) (i) A particle is confined in a one-dimensional potential well defined by

$$V_x = \begin{cases} 0, & 0 < x < a \\ \infty, & x \leq 0 \text{ and } x \geq a \end{cases}$$

Obtain the energy eigenvalues and the normalized eigenfunction of the system.

- (ii) Explain the 'spin-orbit coupling' of atomic electron and the consequent doubling of spectral lines with the necessary expressions. 5+5
- (d) (i) Using the vector atom model, determine the possible terms corresponding to the principal quantum number  $n = 3$ , and compute the angle between  $\vec{l}$  and  $\vec{s}$  vectors for the term  ${}^2D_{5/2}$ .
- (ii) Consider two electrons:  $l_1 = 3, s_1 = \frac{1}{2}; l_2 = 1, s_2 = \frac{1}{2}$ . Find the  $J$  values assuming  $J$ - $J$  coupling. (3+3)+4